

Cognitive Behavioral Therapy (CBT) and Virtual Reality (VR) for Enhanced Therapeutic Outcomes: Exposure Therapy



- ▶ Cognitive Behavioral Therapy (CBT) is an evidence-based psychotherapeutic approach that addresses dysfunctional emotions, behaviors, and thoughts through a goal-oriented, systematic process. CBT is designed to help individuals identify and challenge unhelpful cognitive distortions (e.g., black-and-white thinking, overgeneralization, catastrophizing) and develop healthier thought patterns and behaviors. The efficacy of CBT has been extensively documented in the treatment of various mental health disorders, including anxiety, depression, post-traumatic stress disorder (PTSD), and obsessive-compulsive disorder (OCD) (Tolin, 2010; Hofmann et al., 2012; David et al., 2018).
- ▶ Exposure therapy is a specific type of CBT that involves the gradual and repeated exposure of patients to feared objects, activities, or situations in a safe and controlled environment. The aim is to reduce the fear response through habituation and extinction. Exposure therapy is particularly effective for anxiety disorders, including phobias, social anxiety, and PTSD (Foa et al., 2007; Powers & Emmelkamp, 2008). The process helps patients confront their fears and learn that the feared outcomes are unlikely to occur or are not as catastrophic as anticipated.
- ▶ Virtual Reality (VR) is a technology that creates a simulated environment, providing users with immersive and interactive experiences. By wearing VR headsets, individuals can be transported into realistic virtual environments that can be manipulated to suit various therapeutic needs. This technology has gained attention for its potential to enhance traditional therapeutic methods, including exposure therapy, by providing controlled, repeatable, and safe exposure scenarios (Maples-Keller et al., 2017).
- ▶ Virtual Reality Exposure Therapy (VRET) is an innovative modality of exposure therapy used in the treatment of various psychological disorders, including post-traumatic stress disorder (PTSD) and specific phobias. This technique employs computer-generated virtual environments to simulate anxiety-triggering situations for patients, allowing for controlled and safe exposure to their fears (Rothbaum et al., 1995). Botella et al. (2007) demonstrated the efficacy of VRET in treating phobias, highlighting its potential in clinical settings. Craske et al. (2008) further emphasized the importance of exposure therapy in anxiety treatment, supporting the integration of VRET into therapeutic practices. VRET is based on principles of classical conditioning and gradual exposure, facilitating an immersive experience that can be tailored to address specific emotional responses (Freeman et al., 2017). This approach has been shown to be effective in reducing symptoms and improving patients' quality of life by providing a flexible and replicable therapeutic environment (Huang et al., 2016).



Integration of CBT, VR, and Exposure Therapy

Enhanced Therapeutic Efficacy

- ▶ The integration of VR with CBT and exposure therapy has shown promising results in enhancing therapeutic efficacy. VRET allows for the precise control of exposure scenarios, enabling therapists to tailor the intensity and duration of exposure to the individual needs of each patient. This level of control is often not possible with in vivo (real-life) exposure therapy (Botella et al., 2017). Studies have demonstrated that VRET is as effective, if not more, than traditional exposure therapy in reducing symptoms of anxiety disorders and PTSD (Carl et al., 2019; Morina et al., 2015).
- ▶ For instance, a meta-analysis by Carl et al. (2019) found that VRET resulted in significant reductions in symptoms of anxiety and PTSD, comparable to those achieved through traditional exposure therapy. Additionally, the immersive nature of VR can increase patient engagement and adherence to therapy, which are critical factors in achieving successful therapeutic outcomes (Freeman et al., 2017).

Improved Efficiency

- ▶ The use of VR in CBT also improves the efficiency of therapy sessions. VR allows therapists to recreate specific situations or environments that might be difficult or impractical to replicate in real life. This capability enables more efficient use of therapy time and resources. For example, a therapist can simulate a fear of flying scenario without the need to arrange actual flights or create a virtual social interaction for someone with social anxiety, all within the confines of the therapist's office (Freeman et al., 2017).
- ▶ Furthermore, VR can help remote therapy sessions, making mental health services more accessible to individuals in underserved or remote areas. This accessibility can reduce the overall economic burden on healthcare systems by decreasing the need for long-term treatment and improving public health outcomes (Maples-Keller et al., 2017).

VRET and XRHeath external research

- ▶ Research on the feasibility, efficiency, and effectiveness of VRETm using XRHeath's platform has yielded promising results across a range of psychological conditions.

Fear of Darkness in Children

- ▶ At the University of the Balearic Islands, a study aimed to treat children aged 8 to 12 with fear of darkness using VR environments focused on darkness and storms. The program involved 6 to 8 sessions, each lasting approximately 40 minutes. Results showed significant post-treatment improvements in fear levels, with 66.6% of participants showing satisfactory improvement and 16.6% showing acceptable improvement. This suggests that VR interventions can be effectively administered by non-expert therapists, making them a possible treatment choice for common childhood phobias (Navarro-Oliver, 2017; Servera et al., 2019).

Agoraphobia in Adults

- ▶ The TICSalut Foundation and the Institut de Neuropsiquiatria i Addiccions (INAD) conducted a study to improve agoraphobia treatment using VR. The intervention included one session of psychoeducation and five sessions of cognitive restructuring combined with VRET. Findings proved improvements in patients' autonomy, mobility, and overall activity, with 98% of participants requiring only the scheduled sessions, 87% adhering to the treatment, and 72% expressing satisfaction. This highlights the efficiency and effectiveness of integrating VR into standard psychological treatment protocols (Gelabert, 2017; Gelabert et al., 2018).

Fear of Flying

- ▶ Hofstra University compared remotely conducted VRET with traditional office-based VRET for treating aviophobia. Over six intensive sessions (100 minutes each), including four VR sessions, results indicated similar outcomes in both settings. Participants' psychometric scores and graduation of flight exposures were comparable, although the remote group experienced more technological issues. This supports the viability of remote VRET, which can improve access to treatment for those unable to attend in-person sessions (Cherestal et al., 2018; Tusa et al., 2018).
- ▶ At Hospital del Mar, a case study focused on a woman with a severe fear of flying. The treatment involved one psychoeducation session followed by progressive VR exposure over six sessions, culminating in a real flight. The patient's anxiety reduced by over 50% between the first and last sessions, and she reported significantly lower anxiety during the actual flight. This demonstrates VR's effectiveness in treating specific phobias, comparable to traditional in vivo exposure (Rosado et al., 2018).

Public Speaking Anxiety

- ▶ A study at Universitat Pompeu Fabra aimed to reduce public speaking anxiety (PSA) among university students using VR environments simulating an auditorium with distractors. The experimental group underwent five VR training sessions, while the control group received traditional public speaking training. Anxiety levels, measured via electrodermal activity and self-report, significantly decreased in the experimental group, particularly among male students who initially reported higher anxiety levels. This highlights VR's effectiveness in addressing PSA, with participants also reporting a positive training experience (Pompeu Fabra University).

Childhood Social Anxiety Disorder

- ▶ The University of Central Florida explored the use of VR environments in treating social anxiety disorder in children aged 8 to 12. The study replaced in vivo peer generalization sessions and standard parent-assisted homework with VR practice. Results showed high satisfaction among children, parents, and clinicians, with VR environments viewed as acceptable, feasible, and credible treatment components. This suggests VR can effectively complement traditional therapy for social anxiety disorder, providing ample practice opportunities and ensuring treatment generalizability (Wong, Beidel, & Spitalnick, 2022).
- ▶ These studies underscore the potential of XRHealth's VR platform in enhancing therapeutic outcomes for various psychological conditions. By demonstrating efficacy, efficiency, and feasibility, VR interventions present a significant advancement in mental health treatment, offering accessible, engaging, and effective therapy options.

Scientific Justification

- ▶ The scientific literature supports the integration of VR with CBT and exposure therapy as a method to enhance therapeutic outcomes and efficiency. Studies have shown that VRET can lead to faster and more durable treatment effects compared to traditional methods. A review by Botella et al. (2017) highlighted the efficacy of VRET in treating PTSD, with patients showing significant improvements after fewer sessions compared to traditional exposure therapy.
- ▶ Moreover, a study by Rizzo and Shilling (2017) showed that the first costs associated with implementing VR technology are offset by long-term savings in treatment costs and improved clinical outcomes. This finding suggests a positive return on investment (ROI) for healthcare providers adopting VR-enhanced CBT.

References:

- ▶ Botella, C., Baños, R. M., Villa, H., Perpiñá, C., & García-Palacios, A. (2007). Virtual reality in the treatment of claustrophobic fear: A controlled, multiple-baseline design. *Behavior Therapy*, 31(3), 583-590.
- ▶ Botella, C., Serrano, B., Baños, R. M., & Garcia-Palacios, A. (2017). Virtual reality exposure-based therapy for the treatment of post-traumatic stress disorder: A review of its efficacy, the adequacy of the treatment protocol, and its acceptability. *Neuropsychiatric Disease and Treatment*, 13, 2533-2545. <https://doi.org/10.2147/NDT.S123640>
- ▶ Carl, E., Stein, A. T., Levihn-Coon, A., Pogue, J. R., Rothbaum, B., Emmelkamp, P., Asmundson, G. J. G., Carlbring, P., & Powers, M. B. (2019). Virtual reality exposure therapy for anxiety and related disorders: A meta-analysis of randomized controlled trials. *Journal of Anxiety Disorders*, 61, 27-36. <https://doi.org/10.1016/j.janxdis.2018.08.003>
- ▶ Craske, M. G., Kircanski, K., Zelikowsky, M., Mystkowski, J., Chowdhury, N., & Baker, A. (2008). Optimizing inhibitory learning during exposure therapy. *Behaviour Research and Therapy*, 46(1), 5-27.
- ▶ Cherestal, S., Schare, M. L. & Tusa, B. (2018, November). Remotely conducted versus office-based virtual reality treatment for fear of flying. 52nd Annual Convention of the Association for Behavioral & Cognitive Therapies, Washington, DC, US.
- ▶ Cherestal, S., Schare, M. L., Grimaldi, S., Costello, K., & DeJesus, C. (2021). Remotely conducted versus office-based virtual reality treatment for aviophobia: Questions of feasibility and accessibility. *Translational Issues in Psychological Science*, 7(3), 218–228. <https://doi.org/10.1037/tps0000303>
- ▶ David, D., Cristea, I., & Hofmann, S. G. (2018). Why cognitive behavioral therapy is the current gold standard of psychotherapy. *Frontiers in Psychiatry*, 9(JAN), 6–8. <https://doi.org/10.3389/fpsy.2018.00004>
- ▶ Foa, E. B., Hembree, E. A., Cahill, S. P., Rauch, S. A. M., Riggs, D. S., Feeny, N. C., & Yadin, E. (2007). Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: Outcome at academic and community clinics. *Journal of Consulting and Clinical Psychology*, 73(5), 953-964. <https://doi.org/10.1037/0022-006X.75.5.903>
- ▶ Freeman, D., Haselton, P., Freeman, J., Spanlang, B., Kishore, S., Albery, E., Denne, M., Brown, P., Slater, M., & Nickless, A. (2017). Automated psychological therapy using immersive virtual reality for treatment of fear of heights: A single-blind, parallel-group, randomised controlled trial. *The Lancet Psychiatry*, 5(1), 17-25. [https://doi.org/10.1016/S2215-0366\(17\)30406-7](https://doi.org/10.1016/S2215-0366(17)30406-7)

- Gelabert, J. M. (2017, December). Agora VR. [Tecnologia 4.0. (experiències + debat)]. TIC, Salut i Social, Mataró, Spain.
- Gelabert-Mir, J. M., & Giner-Bartolomé, C. (2018, September). Intervenció psicològica de l'agorafòbia mitjançant realitat virtual: Avaluació de paràmetres d'eficiència. 9es Jornades R+D+I TIC Salut i Social, Vic, Spain.
- Hofmann, S. G., Asnaani, A., Vonk, I. J., Sawyer, A. T., & Fang, A. (2012). The Efficacy of Cognitive Behavioral Therapy: A Review of Meta-analyses. *Cognitive Therapy and Research*, 36(5), 427-440. <https://doi.org/10.1007/s10608-012-9476-1>
- Huang, H., Yang, C., & Liu, J. (2016). The effectiveness of virtual reality exposure therapy on post-traumatic stress disorder: A meta-analysis. *Journal of Anxiety Disorders*, 41, 13-21.
- Maples-Keller, J. L., Bunnell, B. E., Kim, S. J., & Rothbaum, B. O. (2017). The use of virtual reality technology in the treatment of anxiety and other psychiatric disorders. *Harvard Review of Psychiatry*, 25(3), 103-113. <https://doi.org/10.1097/HRP.0000000000000138>
- Morina, N., Ijntema, H., Meyerbröker, K., & Emmelkamp, P. M. (2015). Can virtual reality exposure therapy gains be generalized to real-life? A meta-analysis of studies applying behavioral assessments. *Behaviour Research and Therapy*, 74, 18-24. <https://doi.org/10.1016/j.brat.2015.08.010>
- Rizzo, A. S., & Shilling, R. (2017). Clinical Virtual Reality tools to advance the prevention, assessment, and treatment of PTSD. *European Journal of Psychotraumatology*, 8(sup5), 1414560. <https://doi.org/10.1080/20008198.2017.1414560>
- Rodero, E., & Larrea, O. (2022). Virtual reality with distractors to overcome public speaking anxiety in university students. [Realidad virtual con distractores para superar el miedo a hablar en público en universitarios]. *Comunicar*, 72, 87-99. <https://doi.org/10.3916/C72-2022-07>
- Rosado, S., Aliart, X., Chamorro, J., & Aceña, R. (2018, April). Intervención de enfermería ante las fobias mediante Realidad Virtual. Poster at Congreso Nacional de Enfermería de Salud Mental, Ávila, Spain.
- Rothbaum, B. O., Hodges, L., Kooper, R., & Opdyke, D. (1995). Virtual reality exposure therapy for PTSD Vietnam veterans. *Journal of Clinical Psychiatry*, 56(4), 138-144.
- Servera, M., Sáez, B., & Gelabert-Mir, J. M. (2019). Feasibility of a virtual reality program to treat children with fear of darkness with nonexpert therapists. *Revista de Psicología Clínica con Niños y Adolescentes*, 7(2), 16-21. <https://doi.org/10.21134/rpcna.2020.07.2.2>.
- Tolin, D. F. (2010). Is cognitive-behavioral therapy more effective than other therapies? A meta-analytic review. *Clinical Psychology Review*, 30(6), 710-720. <https://doi.org/10.1016/j.cpr.2010.05.003>
- Tusa, B., Schare, M. L., Argento, V., Sanderson, W. & Buqo, T. (2018, November). The Effect of Remote virtual reality exposure therapy on presence and related technological issues. 52nd Annual Convention of the Association for Behavioral & Cognitive Therapies, Washington, DC, US.
- Wong Sarver, N., Beidel, D. C., & Spitalnick, J. S. (2014). The Feasibility and Acceptability of Virtual Environments in the Treatment of Childhood Social Anxiety Disorder. *Journal of Clinical Child & Adolescent Psychology*, 43(1), 63-73. <https://doi.org/10.1080/15374416.2013.843461>

Thank You.



Follow Us



www.xr.health